

NOVEL IMMUNO-STIMULATING POLYSACCHARIDE SUBSTANCE FROM
Phellinus spp. STRAIN AND USE THEREOF

EXPRESS MAIL CERTIFICATE

Date 12/28/98 Serial No. 5445425920

I hereby certify that on the date indicated above I deposited this paper or fee with the U.S. Postal Service & that it was addressed for delivery to the Commissioner of Patents & Trademarks, Washington, DC 20231 by "Express Mail Post Office to Addressee" service.

Name (Print) D B Peck

Signature D B Peck

BACKGROUND OF THE INVENTION

1. Field of the invention

The present invention relates to a strain belonging to genus *Phellinus*, a polysaccharide substance produced from the strain, and use of the polysaccharide substance. More particularly, the present invention relates to a novel *Phellinus* spp. strain, a novel polysaccharide substance produced therefrom, which shows potent immuno-stimulating activity, and use of the polysaccharide substance in the prophylaxis and treatment of immunity-related diseases, such as cancers and AIDS, and in their mechanism study.

2. Description of the Prior Art

Mushrooms, a kind of higher fungi, are heterotrophic; they take in organic nutrients from the outside through microscopically structured hyphae. In the biosphere, they play a role as a decomposer which degrade complex organic materials into simple inorganic forms and finally return them into nature. Of fungi, most typical mushrooms belong to Basidiomycetes which characteristically form basidia which each bears on its surface four exogenous basidiospores upon coming to maturity. The mushrooms are further classified into Hymenomycetes which

5/1/01 82853600842US

D B Peck D B Peck

actively release spores from exposed hymenia. Representative of Hymenomycetes are Aphyllophorales and Agaricales. *Ramaria botrytis*, *Hymenochaete yasudai* Imaz., *Climacodon septentrionalis*, *Coriolus versicolor*, and *Canoderma applanatum* fall under the category of Aphyllophorales and *Pleurotus ostreatus*, *Lentinus edodes*, *Schizophyllum commune*, *Tricholoma matsutake*, *Amanita muscaria*, and *Coprinus comatus* are placed within the category of Agaricales.

From times past, it has been known in Oriental herb medicine that many mushrooms belonging to Aphyllophorales, a kind of Basidiomycetes, are therapeutically active against various diseases, especially, tumors. In Korea, they have been transmitted from generation to generation as herb medicines or folk medicines and, among the mushrooms, particularly, *Phellinus linteus*, belonging to the family Polyporaceae, has been regarded as an extremely precious medicine. *Phellinus linteus*, however, is a very rare species in nature, so that its fruiting body is very difficult to obtain. Further, isolation and artificial culture of the mycelium of *Phellinus linteus* are regarded almost impossible. Accordingly, advantage has not been actively taken of *Phellinus linteus* despite its being recognized as a therapeutically very effective agent against tumors.

For treating cancers, various methods including chemotherapy, radiotherapy and surgery are conventionally used. Effective as these therapies are for solid cancer, they are

insufficient for the perfect conquest of caners. Particularly, they are helpless for metastatic tumors.

Usually, chemotherapy using anticancer agents, such as adriamycin, and radiotherapy cause serious side-effects. In order to attenuate the side-effects, such an anticancer agent would be administrated at a reduced dose. In this case, however, the therapeutic effect of the chemical becomes poor. What is worse, cancer metastasis cannot be prevented even after taking chemotherapy in treating cancers. Thus, conventional chemotherapy is effective against the growth of cancers to an extent, but cannot completely cure cancers.

An alternative method is to utilize immunosuppressives, known as immunotherapy. Representative examples of the immunosuppressives are cytokines, such as interleukine-2, tumor necrosis factors, etc. Interleukine 2 shows superior therapeutic effects, particularly for cancer metastasis. When being applied for the human body, interleukine-2 is required to be used at such a large amount as to cause serious side-effects. Recent research on novel immuno-stimulating substances which are non-cytokines and show excellent activity against cancer without side-effect, has resulted in the development of Lentinan and OK-432.

For last few years, a new technology in which chemotherapy is used in combination with immunotherapy, known as immunochemotherapy, has been gathering strength. For instance, adriamycin and the immnuoenhancements are co-administrated to

the patients suffering from cancer, with the aim of enhancing the anticancer effects while reducing the side effects of the chemical.

No study on the microbiological and genetical characteristics of *Phellinus* spp. strains has been, to our knowledge, reported, thus far. Microorganisms, even if the same species, may be significantly different from each other in the kinds and amounts of the materials they produce as well as may show a large diversity in culture conditions, strain stability, and frequency of genetic variation. The first thing to do in researching the fungi, *Phellinus*, on their valuable products is to select the strains which can produce a large amount of the products and be cultured in broth in addition to being of low genetic variation. *Phellinus linteus* strains are very diverse and modified in morphology as observed with a microscope, so that they are difficult to identify. However, their classification has been practically dependent on the morphological discrimination under a microscope.

In recent, new DNA analysis methods have been developed for research in population genetics and phylogenetics. Over phenotype analysis methods, the DNA analysis methods have an advantage of being able to analyze genotypes on the basis of evolution rate and heredity tendency. Of them, a restriction fragment length polymorphism (hereinafter referred to as "RFLP") technique allows the mutation of base sequence to be inferred so

as to analyze genotypes (Dowling et al. 1990, 1990. Nucleic acid
II: Restriction site analysis. In: Molecular systematics ed by D.
M. Hills and C. Moritz. P250-317. Sinauer Associates Press).

Because nuclear DNA is too large to easily analyze, organelle

5 genomes are usually utilized for DNA analysis (White and
Densmore, 1992 Mitochondrial DNA isolation. In: Molecular genetic
analysis of populations : A practical approach. Ed by A. R.
Hoelzel. P29-58. Oxford university press). Particularly,
advantage is taken of mitochondrial DNA. Generally,
10 mitochondrial DNA is smaller in size and faster in evolutionary
change rate than nuclear DNA and experiences maternal haploid
inheritance, so that mitochondrial DNA is extensively used for
study on the genetic structure and history of a population.
Particularly, the RFLP of mitochondrial DNA is utilized for the
15 study on inter-species mutation (Foster et al., 1988. Estimation
of relatedness between *Phytophthora* species by analysis of
mitochondrial DNA. *Mycologia* 80:466-478)

SUMMARY OF THE INVENTION

20

Through the intensive and thorough research on *Phellinus*
spp. strains utilizing the DNA analysis method, the present
inventors have found that a novel polysaccharide substance which
a *Phellinus* strain produces, has a potent immuno-stimulating
25 activity.

Therefore, it is an object of the present invention to provide a *Phellinus* spp. strain which produces a novel polysaccharide substance of immuno-stimulating activity.

It is another object of the present invention to provide a
5 novel polysaccharide substance which shows a potent immuno-stimulating activity against immune-related diseases.

It is a further object of the present invention to provide use of the polysaccharide substance as an immunotherapy agent or in the study on the basic mechanism of the diseases.

10 For achieving these objects, first, various *Phellinus* spp. strains were cultured and their total DNA were isolated by a SDS-phenol method. Of the total DNA, only mitochondrial DNA is obtained. The mitochondrial DNA was digested with restriction enzymes BamHI, ClaI, EcoRI and PvuII, followed by electrophoresis
15 on an agarose gel. The DNA relatedness among the *Phellinus* strains was analyzed according to the method of Nei and Li (1979). From the analytical data, a novel strain was found which forms new relatedness to other strains. This strain was named *Phellinus linteus* Yoo and deposited in the Korean Collection for
20 Type Cultures, Korean Research Institute of Bioscience and Biotechnology on Nov. 17, 1997 (Deposition No. KCTC 0399BP). An investigation was made on the microbiological features of the KCTC 0399BP strain. A novel polysaccharide substance was isolated and purified from the fruiting bodies and mycelia of the
25 novel strain and tested for immuno-stimulating activity. The

immuno-stimulating polysaccharide substance was analyzed for its sugar units and composition, followed by the structural analysis of carbohydrate component. Similarly, other various *Phellinus* spp. strains were cultured and their fruiting bodies and mycelia
5 were subjected to isolation and purification to give polysaccharide substances which were also of immuno-stimulating activity.

Animal tests were performed. The immuno-stimulating polysaccharide substances were administered to mice into which
10 cutaneous cancer cells were implanted to measure their anticancer activity. Also, when being co-administrated with chemicals, the polysaccharide substances were measured for their change in anticancer activity. The effect of the polysaccharide substance on the growth of human cancer cells was researched. To mice
15 which were intravenously injected with cutaneous cancer cells at the tails, the polysaccharide substance was administered, in order to investigate its influence on cancer metastasis. In the present invention, the anticancer activity of the polysaccharide substance was found to be different from that of anticancer
20 chemicals in mechanism.

BRIEF DESCRIPTION OF THE DRAWINGS

The above and other objects and aspects of the invention
25 will become apparent from the following description of

embodiments with reference to the accompanying drawings in which:

Fig. 1 is a phylogenetic tree showing the relatedness among *Phellinus* spp. strains;

Fig. 2 is a graph showing the culture properties of the mycelia of *Phellinus linteus*;

Fig. 3 is a procedure showing the purification of the polysaccharide substance from *Phellinus* stains;

Fig. 4 is a gas chromatogram for the sugar units of the immuno-stimulating polysaccharide substance;

Fig. 5 is a HW 65F gel chromatogram for the immuno-stimulating polysaccharide substance;

Fig. 6 is a carbon NMR spectrum for the immuno-stimulating polysaccharide substance;

Fig. 7 is a graph showing the anticancer activities against mouse cancer cells of the polysaccharide substance according to administration manner;

Fig. 8a is a graph showing the anticancer activities against mouse cancer cells of a combination of the polysaccharide substance and adriamycin 0.1 mg/kg;

Fig. 8b is a graph showing the anticancer activities against mouse cancer cells of a combination of the polysaccharide substance and adriamycin 0.3 mg/kg;

Fig. 9a is a graph showing the change of the human cancer cells according to the treatment with the polysaccharide substance and adriamycin;

Fig. 9b is a graph showing the final weight of the human cancer cells according to the treatment with the polysaccharide substance and adriamycin;

Fig. 10a is a graph showing a prevention effect of
5 adriamycin on cancer metastasis;

Fig. 10b is a graph showing a prevention effect of a combination of the polysaccharide substance and adriamycin on cancer metastasis; and

Fig. 11 is a graph showing the cytotoxicity of adriamycin
10 and polysaccharide substance.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

13 *Phellinus* strains including the novel *Phellinus linteus*
15 Yoo KCTC 0399BP are each cultured on a potato dextrose agar (PDA). Using an SDS-phenol technique, a total DNA is isolated from the mycelia of each of the cultures, followed by further isolation of mitochondrial DNA from the total DNA. These
mitochondrial DNAs are useful to analyze the relatedness among
20 the strains. For this analysis, the mitochondrial DNAs are digested with restriction enzymes and the DNA fragments digested are separated on an agarose gel by electrophoresis. The
distances which the DNA fragments move on the gel are the data for calculating restriction enzyme fragment relatedness values
25 (F) and sequence divergence values (p). Based on the values, a

novel strain was found with a relatedness and named "*Phellinus linteus* Yoo" and deposited in the Korean Collection for Type Culture, Korean Research Institute of Bioscience and Biotechnology (Deposition No. KCTC 0399BP). A study on the
5 microbiological features of the fruiting body of *Phellinus linteus* Yoo KCTC 0399BP is helpful in proliferating this mushroom.

In order to extract the substance which is thought to be therapeutically effective, the mushroom is cultured and
10 proliferated. A hot-water extraction gives a novel substance which is proved to be a kind of polysaccharides as analyzed by phenol-sulfuric acid and Folin-phenol techniques. In addition, the structural characteristics of this polysaccharide substance can be revealed by an NMR technique.

15 After being administrated to an animal, the polysaccharide substance is tested for immuno-stimulating activity by counting the number and measuring the proliferation of its lymphocytes. A similar procedure to that for *Phellinus linteus* Yoo is carried out for other *Phellinus* strains. That is, strain culture,
20 mycelium extraction and substance isolation are sequentially done. The substances isolated are tested for Immuno-stimulating activity, too. In addition, the polysaccharide substances which are obtained from the fruiting bodies of the *Phellinus* strains are proved to be of immuno-stimulating activity.

25 In order to assay the activity of the polysaccharide

substance against cancer, animal tests are performed. First,
mouse-derived cutaneous cancer cells are implanted into mice to
which the polysaccharide substance and other chemicals are, then,
administrated alone or in combination. In either case, the
5 viabilities of the mice are compared. The activity of the
polysaccharide against human cancer is measured by using the nude
mice into which human cancer cells are implanted. To the nude
mice, the polysaccharide substance and adriamycin are
administrated alone or in combination. In addition, the
10 polysaccharide substance is very effective in preventing cancer
metastasis. For this, mouse cutaneous cancer cells are
intravenously injected into the tails of mice, followed by
administration of the polysaccharide substance and adriamycin. A
sulforhodamin B method is taken to measure viable mouse cutaneous
15 cancer cells and human lung cancer cells which are previously
treated with the polysaccharide substance and adriamycin alone or
in combination.

A better understanding of the present invention may be
obtained through the following examples which are set forth to
20 illustrate, but are not to be construed as the limit of the
present invention. In the following examples, splenocytes taken
from B₆C₃F mice were used for the measurement of the immuno-
stimulating activity of the polysaccharide substances isolated
from *Phellinus linteus* Yoo and other *Phellinus* spp. First, the
25 splenocytes were immunized for 2 days by treatment with the

substances and lipopolysaccharide as a positive control. The antibodies formed were measured by a spectrophotometer and considered as a standard for the immuno-stimulating activity. The effect of the polysaccharide substance on the proliferation of T-cells was measured by a mixed lymphocyte response technique. To detect the proliferation of lymphocytes, nuclear DNA which was newly synthesized in lymphocytes was measured by using ^3H -thymidine.

10 EXAMPLE I: Culture of *Phellinus* Strain

The *Phellinus* spp. strains as given in Table 1, below, were each cultured on a potato dextrose agar (PDA) at a constant temperature of 28 °C for 5-12 days. Following inoculation in 450 mL of a PD broth, each of the strains was grown at 28 °C for 12 days with agitation at 120 rpm.

TABLE 1

Phellinus spp. used and Origins

Strains	Origins
<i>P.linteus</i> WD1222	Japan
<i>P.linteus</i> ATCC 26710	U.S.A
<i>P.linteus</i> KCTC 0399BP	Korea
<i>P.linteus</i> PL5	Korea
<i>P.linteus</i> PL4	Korea
<i>P.linteus</i> PLM	Korea
<i>P.tremulae</i> ATCC 32233	U.S.A
<i>P.johnsonianus</i> ATCC 60051	U.S.A
<i>P.pini</i> ATCC 12240	U.S.A
<i>P.tuberculosis</i> ATCC 13271	U.S.A
<i>P.nigricans</i> FP-71807-S	U.S.A
<i>P.chrysoloma</i> HHB-3585-Sp	U.S.A

EXAMPLE II: Total DNA Isolation

An SDS-phenol method was utilized for the Isolation of total DNA from mycelia. Each of the mycelia cultured in Example I was washed with 20 mM EDTA, harvested by vacuum filtration using a filter paper (Whatman No. 1), stored in a deep-freezer maintained at -70 °C, and subjected to freeze-drying. The freeze-dried mycelia were finely ground and 2 g of each of these powders were added with 20 mL of an extraction buffer (0.2M Tris·HCl, pH 8.5; 0.25M NaCl; 25mM EDTA; 0.5%(w/v) SDS) and then, with 14 mL of

phenol and 6 mL of chloroform. After being mixed by slow agitation, the solution was centrifuged. The resulting supernatant was treated with 15 μ l of RANase A (10 mg/mL) at 37 °C for 30 min. and then, further with 15 μ l of proteinase K (10 mg/mL) at 60 °C for 15 min. Thereafter, the enzyme-treated supernatant was well mixed with an equal volume of a mixture of phenol, chloroform and isoamyl alcohol (25:24:1), followed by centrifugation at 12,000 X g for 30 min. at 4°C. Before repeating the same centrifugation, the supernatant was well mixed with an equal volume of a mixture of chloroform and isoamyl alcohol (24:1). The supernatant thus obtained was mixed with 0.54 volume of isopropanol by vortexing and centrifuged under the same conditions as in above, to give a DNA pellet. This pellet was washed with 70% ethanol and dried in air. The DNA was dissolved in 8 mL of a TE buffer (10mM Tris·HCl, 1mM EDTA, pH 8.0) and stored at -20 °C.

EXAMPLE III: Mitochondrial DNA Isolation

An ultracentrifugation was carried out to isolate mitochondrial DNA from the total DNA obtained from each of the strains in Example II. After being mixed with 8.8 g of cesium chloride, commercially available from Sigma, U.S.A., and 6 μ l of a bisbenzimidazole solution (10 mg/mL), the total DNA solution (8 mL) was ultracentrifuged at 40,000 X g for 40 hours at 20°C.

Following confirmation of two DNA bands under a uv light, only the upper band, that is, a mitochondrial DNA fraction, was recovered with the aid of a 21 gauge syringe needle. The mitochondrial DNA fraction thus obtained was added with an equal
5 volume of CsCl-saturated isopropanol, mixed by weakly vortexing and centrifuged at 1,200 X g at 4 °C. This washing step was repeated 6-7 times so as to completely remove the bisbenzimidide from the mitochondrial DNA fraction. Removal of the cesium chloride was accomplished by adding 70% ethanol three times in
10 volume than that of the mitochondrial DNA fraction to the mitochondrial DNA and centrifuging at a speed of 100 X g at 4 °C for 10 min. This washing step was repeated once more to give bisbenzimidide- and Cs-free mitochondrial DNA. After being dried in air, the mitochondrial DNA was dissolved in a TE buffer (pH
15 8.0) and quantitated at 260 nm with a spectrophotometer, such as that sold by Beckman, U.S.A under the tradename of "DU-64 Spectrometer". The mitochondrial DNA was stored at -20 °C.

20 EXAMPLE IV: Digestion of the Mitochondrial DNAs with Restriction Enzymes

The restriction enzymes used in the example as well as the buffers therefor were purchased from Boehringer Mannheim (Germany) and are given as listed in Table 2, below, along with
25 their recognition sites. Each of the mitochondrial DNA samples

of the *Phellinus* strains, obtained in Example III, had a concentration of 1 μ g per 10 μ l of buffer and digested at 37 °C for 3 hours.

TABLE 2

Restriction Enzymes and Their Recognition Sites

Restriction Enz.	Recognition Sites
BamHI	G↓GATCC
ClaI	AT↓CGAT
EcoRI	G↓AATTC
PvuII	CAG↓CTG

EXAMPLE V: Electrophoresis on Agarose Gel

The DNA samples which had been treated with the restriction enzymes in Example VI, were mixed with a gel loading buffer (0.25% bromophenol blue, 0.25% xylene cyanol FF, 30% glycerol) and electrophoresed on 1 % agarose gel (agarose MP), sold by Boeringer Mannheim, Germany, for 4 hours in a horizontal electrophesing analyzer, commercially available from BRL, U.S.A., using a TAE buffer (40mM Tris-acetate, 1mM EDTA, pH 8.0) as a running buffer by applying 50 V across the analyzer. After running the DNA samples, the agarose gel was immersed in an ethidium bromide buffer (0.5 μ g/mL in water) to easily detect the DNAs on a UV lamp. The mitochondrial DNA sizes which were

determined by the electrophoresis are given as shown in Table 3, below.

TABLE 3

Sizes of Mitochondrial DNA Isolated from *Phellinus* spp.

Strains	DNA Size
<i>P.linteus</i> WD1222	58Kb
<i>P.linteus</i> ATCC 26710	59Kb
<i>P.linteus</i> KCTC 0399BP	61Kb
<i>P.linteus</i> PL5	61Kb
<i>P.linteus</i> PL4	73Kb
<i>P.linteus</i> PLM	63Kb
<i>P.tremulae</i> ATCC 32233	77Kb
<i>P.johnsonianus</i> ATCC 60051	76Kb
<i>P.pini</i> ATCC 12240	76Kb
<i>P.tuberculosis</i> ATCC 13271	57Kb
<i>P.nigricans</i> FP-71807-S	58Kb
<i>P.chrysoloma</i> HHB-3585-Sp	60Kb

EXAMPLE VI: Restriction Fragment Length Polymorphism (RFLP)
Analysis

The analysis for the DNA relatedness between *Phellinus* strains was carried out according to Nei and Li method. Assuming that the DNA fragments were identical to each other if they moved

the same distance when they were electrophoresed on agarose gel as in Example V, restriction enzyme fragment relatedness values (F) and sequence divergence values (p) were obtained. From the P values, the relatedness between *Phellinus linteus* Yoo (KCTC 0399BP) strain and other *Phellinus* spp. strains was analyzed by an Unweighted Pair-Group Method, arithmetic average (UPGMA) Clustering technique. The P and F values were calculated on the basis of the following equations, respectively:

$$F = 2n_{xy} / (n_x + n_y)$$

wherein F shows a relatedness between two *Phellinus* spp., n_{xy} is the number of fragments in common therebetween, and n_x and n_y are each the total number of the fragments displayed by isolates x and y .

$$p = -(\ln F) / r$$

wherein p is an estimate of the proportion of nucleotide base substitution per nucleotide position and r is the number of the nucleotide base pairs for restriction endonuclease recognition site.

The F values obtained when treating with restriction enzyme BamHI are listed, together with their corresponding p values, in Table 4, the F values when treating with restriction enzyme ClaI,

together with their corresponding p values in Table 5, the F values when treating with restriction enzyme EcoRI, together with their corresponding p values in Table 6, and the F values when treating with restriction enzyme PvuII, along with their corresponding p values in Table 7. Table 8 contains the proportions of the fragments in common and the arithmetic average distances corresponding to the p values listed over the diagonal line extending from the left above to the right bottom in each of the Tables 4-7. A phylogenetic tree was inferred from these results and is given as shown in Fig. 1, strongly suggesting that the strain of the present invention be newly differentiated from *Phellinus* strains with a significant relatedness.

TABLE 4

F and P values from PRLP of the BamHI-Digested Mitochondrial DNA

	WD 1222	ATCC 26710	KCTC 0399BP	PL5	PL4	PLM	ATCC 32233	ATCC 60051	ATCC 12240	ATCC 13271	FP- 71807- S	HHB- 3585- Sp
5	WD1222	-						0.1921	0.4142		0.3465	0.2310
	ATCC 26710	NC	-	0.0257	0.2570	0.3120	0.2841	0.2310		0.2841	0.2203	
	KCTC 0399BP	NC	0.8571	-	0.0000	0.3120	0.2841	0.2310		0.2841	0.3358	
10	PL5	NC	0.8571	1.0000	-	0.3120	0.2841	0.2310		0.2841	0.3358	
	PL4	NC	0.1538	0.1538	0.1538	-		0.3358			0.3243	0.3120
	PLM	NC	0.1818	0.1818	0.1818	NC	-					0.2841
	ATCC 32233	NC	0.2500	0.2500	0.2500	0.1333	NC	-	0.3753	0.2987	0.1735	0.3465
15	ATCC 60051	0.3158	NC	NC	NC	NC	NC	0.1052	-	0.4209	0.3666	0.2411
	ATCC 12240	0.0833	0.1818	0.1818	0.1818	NC	NC	0.1666	0.0800	-	0.2915	0.3996
	ATCC 13271	NC	0.2666	0.1313	0.1333	0.1428	NC	0.3530	0.1111	0.1739	-	0.2203
20	FP- 71807- S	0.1250	NC	NC	NC	NC	NC	NC	0.2353	0.0909	0.2666	-
25	HHB- 3585- SP	0.1250	NC	NC	NC	0.1538	0.1818	0.1250	0.1176	0.2727	NC	NC

TABLE 5

F and P values from PRLP of the ClaI-Digested Mitochondrial DNA

	WD 1222	ATCC 26710	KCTC 0399BP	PL5	PL4	PLM	ATCC 32233	ATCC 60051	ATCC 12240	ATCC 13271	FP- 71807-S	HHB- 3585-Sp	
5	WD1222	-	0.1155	0.1351	0.1527	0.2841	0.2987	0.3243	0.0372	0.1412		0.2310	0.2507
	ATCC 26710	0.5000	-	0.0196	0.0851	0.1686	0.1831	0.2088	0.1527			0.1155	0.2570
	KCTC 0399BP	0.4444	0.8888	-	0.0531	0.1831	0.1288	0.2203	0.1686			0.1351	0.2682
10	PL5	0.4000	0.6000	0.7272	-	0.1964	0.1412	0.3465	0.1831			0.1527	0.2841
	PL4	0.1818	0.3636	0.3333	0.3077	-	0.1047	0.2411	0.1964	0.1736	0.1256	0.1686	0.2987
	PLM	0.1666	0.3333	0.4615	0.4285	0.5333	-	0.3666	0.3243	0.1831	0.1831	0.1831	0.3120
15	ATCC 32233	0.1428	0.2587	0.2666	0.1250	0.2353	0.1111	-	0.2310	0.2006	0.2006	0.2088	0.2203
	ATCC 60051	0.8000	0.4000	0.3636	0.3333	0.3077	0.1428	0.2500	-		0.3465	0.1682	0.2841
	ATCC 12240	0.4285	NC	NC	NC	0.3529	0.3333	0.3000	NC	-	0.2682	0.3243	0.2203
20	ATCC 13271	NC	NC	NC	NC	0.4706	0.3333	0.3000	0.1250	0.2000	-	0.3243	
	FP- 71807-S												
		0.2500	0.5000	0.4444	0.4000	0.3636	0.3333	0.2857	0.2000	0.1428	0.1482	-	0.2507
25	HHB- 3585-Sp												
		0.2222	0.2222	0.2000	0.1818	0.1666	0.1538	0.2666	0.1818	0.2666	NC	0.2222	-

TABLE 6

F and P values from PRLP of the EcoRI-Digested Mitochondrial DNA

	WD	ATCC	KCTC	PL5	PL4	PLM	ATCC	ATCC	ATCC	ATCC	FP-	HHB-	
	1222	26710	0399BP				32233	60051	12240	13271	71807-S	3585-SP	
5	WD1222	-	0.3837	0.1921	0.2006	0.2764	0.2597	0.3302	0.0175	0.4339	0.1899	0.4071	0.2987
	ATCC 26710	0.1000	-	0.1442	0.2006	0.1608	0.2597	0.3302	0.3837	0.3183	0.3054	0.4071	0.4142
	KCTC 0399BP	0.3158	0.4210	-	0.0509	0.2006	0.2507	0.2567	0.4275	0.3120	0.2310	0.2165	0.2915
	PL5	0.3000	0.3000	0.7368	-	0.2764	0.2597	0.1470	0.2682	0.3183	0.2378	0.2240	0.2987
10	PL4	0.1904	0.3809	0.3000	0.1904	-	0.3837	0.2682	0.2764	0.4399	0.3120	0.2987	0.3054
	PLM	0.2105	0.2105	0.2222	0.2105	0.1000	-	0.3243	0.2597	0.3120	0.4142		0.1964
	ATCC 32233	0.1397	0.1379	0.2143	0.4138	0.2000	0.1428	-	0.3302	0.3662	0.1256	0.3465	0.2362
	ATCC 60051	0.9000	0.1000	0.0769	0.2000	0.1904	0.2105	0.1379	-	0.4339	0.2738	0.2915	0.4142
15	ATCC 12240	0.0740	0.1481	0.1538	0.1481	0.0714	0.1538	0.1111	0.0740	-	0.3465	0.2203	0.2737
	ATCC 13271	0.3200	0.1600	0.2500	0.2400	0.1538	0.0833	0.4705	0.2400	0.1250	-	0.2568	0.2915
	FP- 71807-S	0.0869	0.0869	0.2727	0.2680	0.1666	NC	0.1250	0.1739	0.2666	0.2142	-	0.4339
	HHB- 3585-SP	0.1666	0.0833	0.1739	0.1666	0.1600	0.3077	0.2424	0.0833	0.1935	0.1379	0.0740	-
20													
25													

TABLE 7

F and P values from PRLP of the PvuII-Digested Mitochondrial DNA

	WD	ATCC	KCTC	PL5	PL4	PLM	ATCC	ATCC	ATCC	ATCC	FP-	HHB-	
	1222	26710	0399BP				32233	60051	12240	13271	71807-	3585-	
											S	SP	
5	WD1222	-	0.2507			0.2682	0.2411	0.0851	0.3567	0.2987	0.3358	0.3465	
	ATCC 26710	0.2222	-	0.1155	0.1155	0.2987	0.2507	0.3465	0.3465		0.3243		
	KCTC 0399BP	NC	0.5000	-	0.0479				0.3465		0.3243		
	PL5	NC	0.5000	0.7500	-				0.3465				
10	PL4	NC	0.1666	NC	NC	-	0.1968	0.2682		0.3358			
	PLM	0.2000	0.2222	NC	NC	0.3070	-	0.3567	0.2682				
	ATCC 32233	0.2353	0.1250	NC	NC	0.2000	0.1176	-		0.2310	0.3996	0.4071	
15	ATCC 60051	0.6000	NC	NC	NC	NC	0.2000	NC	-	0.567	0.2987	0.3358	0.3465
	ATCC 12240	0.1176	0.1250	0.1250	0.1250	NC	NC	0.2500	0.1176	-	0.2597	0.1010	0.1760
	ATCC 13271	0.1666	NC	NC	NC	0.1333	NC	NC	0.1666	0.2105	-	0.1256	0.2507
20	FP- 71807-	0.1333	0.1428	0.1428	NC	NC	NC	0.0909	0.1333	0.5454	0.4706	-	0.2088
	S												
25	HHB- 3585-	0.1250	NC	NC	NC	NC	NC	0.0869	0.1250	0.3478	0.2222	0.2857	-
	SP												

TABLE 8

Proportions of fragments in common & arithmetic average distances corresponding to the p values

		WD 1222	ATCC 26710	KCTC 0399BP	PL5	PL4	PLM	ATCC 32233	ATCC 60051	ATCC 12240	ATCC 13271	FP- 71807- S	HHB- 3585- SP
5	WD1222	-	4/53	5/53	5/55	3/60	4/54	5/78	19/59	6/82	5/68	5/62	6/65
	ATCC 26710	0.2499	-	16/50	14/52	8/60	6/51	7/75	3/56	5/79	4/65	4/59	2/62
	KCTC 0399BP	0.1636	0.0762	-	21/52	6/57	6/51	7/75	3/56	5/79	4/65	6/59	3/62
10	PL5	0.1766	0.1645	0.0379	-	5/59	6/51	9/77	4/58	5/81	4/67	5/61	3/64
	PL4	0.2802	0.2350	0.2393	0.2616	-	7/58	8/82	4/63	4/86	8/72	4/66	4/69
	PLM	0.2755	0.2444	0.2212	0.2283	0.2284	-	4/76	4/57	5/80	4/66	2/60	6/63
15	ATCC 32233	0.2985	0.2791	0.2360	0.2415	0.2783	0.3490	-	5/81	10/104	14/90	5/74	8/87
	ATCC 60051	0.0823	0.2682	0.2980	0.2256	0.2364	0.2840	0.3121	-	3/85	6/71	6/65	4/68
20	ATCC 12240	0.3365	0.3163	0.3142	0.3163	0.3067	0.2475	0.2741	0.4038	-	6/94	12/88	12/91
	ATCC 13271	0.2443	0.2628	0.2834	0.2868	0.2744	0.2986	0.1665	0.3123	0.2914	-	10/74	4/77
	FP- 71807- S	0.3301	0.2823	0.2253	0.1883	0.2336	0.1831	0.3183	0.2591	0.2613	0.2317	-	5/71
25	HHB- 3585- SP	0.3817	0.3324	0.2498	0.2914	0.3053	0.2641	0.3025	0.3503	0.2216	0.2711	0.2978	-
30													

EXAMPLE VII : Microbiological Features of *Phellinus linteus* Yoo
Strain KCTC 0399BP

The strain, *Phellinus linteus* Yoo KCTC 0399BP, which was
5 identified as a novel strain newly differentiated from *Phellinus*
with a great relatedness, was studied on its microbiological
features. Its fruiting body, germless and perennial, had a woody
part with a semi-circular pileus about 10 cm wide and about 4 cm
thick. At the time of being picked, this mushroom had a dark
10 brown surface, but, with the lapse of time, the surface came to
have many fissures thereon and the color became deeper to black
brown while its hymenial layer was changed in color from fresh
red to yellowish brown. The tube of the hymenial layer had a
multilayer structure with a fine pore rounded. Its spore 3-4 cm
15 in length was subglobose and light yellowish brown-colored.
Also, it had many setae and a chlamido in a dimension of 15-30 X
8-10 μm . The strain of the present invention was found to be
identical in almost all aspects to the *Phellinus linteus* written
in "Colored Illustrations of Mushrooms of Japan", p 189, printed
20 by Hoikusha Imazeki, edited by Imazeki, R. and Hongo, T., and
named *Phellinus linteus* Yoo. We deposited this novel strain to
Korean Collection for Type Cultures, Korean Research Institute of
Bioscience and Biotechnology, on Nov. 17, 1997, and received a
deposition No. KCTC 0399BP.

EXAMPLE VIII : Culture of *Phellinus linteus* Yoo KCTC 0399BP

A culture broth containing starch 50 g, peptone 10 g and yeast extract 10 g per distilled water 1 L, supplemented with 5 KH_2PO_4 0.8 g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5 g and CaCl_2 0.3 g, was adjusted to pH 6.0 and autoclaved. The mycelia of *Phellinus linteus* Yoo KCTC 0399BP were inoculated in 3.5 liters of the culture broth in a 5 liter fermentation bath and cultured at 28 °C for 11 days while aerating the broth at 2 vvm and rotating the bath at 160 rpm. At 10 5 days after culture, the highest yield for mycelia (30 g/l) was obtained, as shown in Fig.2. From the mycelia cultured for 5 days, polysaccharide substances were hot water-extracted at a yield of 0.873 % per weight of dried mycelia.

15 TABLE 9

Production Yields for Mycelia of *P. linteus* and Polysaccharide

Items	Mycelia of 5 day-Cultured <i>P. linteus</i>
Dried Mycelia Wt.(g/L)	30
Dried Crude Polysaccharide Wt.	262
Crude Polysaccharide Yield (%)	0.873

20 EXAMPLE IX : Extraction of Mycelium Culture of *Phellinus linteus*
25 Yoo strain

An extraction and purification process was done on a mycelium culture of *Phellinus linteus* Yoo strain, in order to obtain crude immuno-stimulating substances. First, 280 g of the mycelium culture obtained in Example VIII was subjected to hot water extraction, followed by concentration to give 3 g of a hot-water extract. This extract was added with ethanol to a final concentration of 80 %, allowed to stand at 4 °C for 24 hours, and centrifuged. The pellet was dissolved in water and then, dialyzed 10 °C for 72 hours using a cellulose tube, such as that sold by Nippon Junko Jun Yaku Seihing, while the dialyzing solution was refreshed every 12 hours. The solution in the cellulose tube was freeze-dried to give a crude extract of 2.45 g.

15 EXAMPLE X : Isolation of Pure Immuno-Stimulating Substance

Purification was achieved by carrying out ion exchange chromatography and gel filtration chromatography successively.

First, the crude extract obtained in Example IX was dissolved in a 5 mM sodium phosphate buffer (pH 7.2), loaded on a DEAE-cellulose column and washed with the same buffer. A solution of NaCl in the same buffer having a concentration gradient from 0 to 1M was used as an eluting buffer. It was flowed through the column at a rate of 5 mL/min. and the eluate was collected at 15 mL per test tube. The eluted fractions were

measured with respect to sugar content by a phenol-sulfuric method and with respect to protein content by Bradford's method. The fractions containing sugar were applied to the top of a column consisting of porous beads, such as that sold under the trade name of Toyopearl HW 65F, and further purified by gel filtration chromatography eluting with water, to afford a pure active fraction which was homogeneous in molecular weight and properties. Its sugar and protein content were measured by the same methods as in above. The pure immuno-stimulating substance thus obtained was named "PL".

EXAMPLE XI : Immuno-Stimulating Activity of the Polysaccharide Substance

The polysaccharide substance derived from Basidiomycetes are known to have a positive influence on the proliferation or activity of lymphocytes, thereby increasing immunity against external factors and even cancer cells. This function of the polysaccharide substance is not attributed to direct cytotoxicity but results from immunological enhancement, so that they advantageously produce little side-effects in vivo. That is, the polysaccharide substances of Basidiomycetes induce an enhancement in the cytotoxic function and antibody formation of the lymphocytes, B-cells and T-cells, thereby showing a potent anticancer immunoactivity. Therefore, the polysaccharide

substances can be indirectly proved to be of anticancer immunoactivity by counting the number of anti-forming lymphocytes and measuring the proliferation of T-cells.

In this example, B-cells were counted by use of an antibody-forming cell method and the proliferation of T-cells was measured by a mixed lymphocyte response technique. In addition, in order to detect the effect of the polysaccharide substance on the proliferation of lymphocytes, the nuclear DNA which was newly synthesized was measured by using ^3H -thymidine. The results are given in Table 10, below.

TABLE 10

Immuno-Stimulating Activity of various Substances

Substance	Lymphocyte Proliferation effect (DPM X 10^4)	T-Cell Activity (DPM X 10^4)	B-Cell Antibody- forming (Abs 576nm)
<i>P.linleus</i> Yoo	8.4 ± 0.5	2.5 ± 0.1	0.66 ± 0.01
LPS	-	-	0.81 ± 0.23
Control	1.6 ± 0.5	0.5 ± 0.05	0.00 ± 0.00

* each substance was used at a concentration of $10 \mu\text{g/mL}$.

As apparent from the data of Table 10, the polysaccharide substance extracted from the novel *Phellinus* strain of the present invention are superior to the control in lymphocyte proliferation and T-cell activity, both, by the factor of about

5-fold. The polysaccharide substance of the present invention show a high B-cell antibody forming power similar to that of lipopolysaccharide. Lipopolysaccharide is a very potent immunosuppressive agent but cannot be freely used owing to its
5 serious side-effects in vivo.

EXAMPLE XII : Sugar Unit and Composition of the Immuno-Stimulating Polysaccharide Substance

10 The polysaccharide substance obtained from Example X was subjected to gas chromatography to analyze its sugar units (Fig. 4). A phenol-sulfuric acid method and a Folin-phenol method were taken to measure its carbohydrate and protein components, respectively. The analysis carried out in an apparatus,
15 commercially available from Hewlett Packard, identified as "HP 5890 GC", provided with a fused silica capillary column sp-2330 (0.32mm X 30m) under the following conditions. The column was first maintained at 200 °C for 2 min. and heated at an elevation rate of 4 °C per min. to 250 °C which was, then, maintained for 2
20 min. A detector and a injector, provided to the apparatus, were maintained at 260 °C and 250 °C, respectively. 2 mg was taken from each of the extract fractions obtained in Example XI, hydrolyzed with 2N TFA, reduced with NaBH₄ and acetylated with anhydrous acetic acid to give aditol acetate for use in gas
25 chromatography. The results are given in Table 11, below.

TABLE 11

Sugar Units and Composition of the Immuno-Active Fraction

Polysaccharide Extract	Carbo- hydrate (%)	Protein (%)	Sugar Units (mole %)		
			Glucose	Monnose	Gatactose
PL	82.7	17.3	78.6	18.0	3.4

EXAMPLE XIII : Structural Analysis of the Immuno-Stimulating Polysaccharide Substance

The Immuno-stimulating polysaccharide substance was subjected to structural analysis.

First, its purity was confirmed by use of 65F gel column chromatography. Fig. 5 is the results of this chromatography.

As seen, a single symmetric peak appeared for each carbohydrate and protein component, demonstrating that PL was pure.

From a carbon nuclear magnetic resonance spectrum (Fig. 6), PL was identified as a glactomannoglucan in which glucose units were joined by $\alpha(1-4)$ and $\beta(1-6)$ linkages to form a long chain and mannose and galactoses were bonded as branches to the long chain.

This structural analysis accorded with the composition result, revealed that PL was a novel polysaccharide quite different in structure from β -glucan.

EXAMPLE XIV : Extraction of Polysaccharide Substances from

Mycelia of *Phellinus* spp. Their Purification and Immuno-Stimulating Activity

Phellinus spp. strains, including *Phellinus igniarius*, *P. baumi*, *P. pini*, *P. gilvus* and *P. nigricans*, were cultured to obtain their mycelia. From them, polysaccharide substances were extracted and purified, and then, used to induce an enhancement in the cytotoxic function and antibody formation of the lymphocytes, B-cells and T-cells, as in Examples VIII to XI. Their anticancer immunoactivities were indirectly evaluated by counting the number of anti-forming lymphocytes and measuring the proliferation of T-cells. Obtained were similar results to those of Example XI.

EXAMPLE XV : Extraction of Polysaccharide Substances from Fruiting bodies of *Phellinus* spp. Their Purification and Immuno-Stimulating Activity

Phellinus spp. strains, including *Phellinus igniarius*, *P. baumi*, *P. pini*, *P. gilvus* and *P. nigricans*, were cultured to obtain their fruiting bodies. From them, polysaccharide substances were extracted and purified, and then, used to induce an enhancement in the cytotoxic function and antibody formation of the lymphocytes, B-cells and T-cells, as in Examples VIII to XI. Their anticancer immunoactivities were indirectly evaluated

by counting the number of anti-forming lymphocytes and measuring the proliferation of T-cells. Obtained were similar results to those of Example XI.

5 EXAMPLE XVI : Therapeutic Activity of the Polysaccharide
Substance against Cancer Cells According to Administration Time

10 B16F10 cutaneous cancer cells, derived from mouse, were
utilized for the measurement of the anticancer activity of the
polysaccharide substance. B16F10 cutaneous cells were cultured
in RPMI 1640 media supplemented with 10 % serum and subcultured
twice every week. As many as 100,000 cells of B16F10 per mouse
were implanted in the peritoneal cavity. The mice implanted were
recorded with respect to viability for 30 days. The recording
15 was also performed for the mice which were peritoneally
administrated with the polysaccharide substance from a week
before the cancer cell implantation to 12 days after the
implantation (pre-treatment) and for the mice which were
peritoneally administrated with the polysaccharide substance for
20 12 days after the implantation (post-treatment). The results are
given as shown in Fig. 7 and summarized in Table 12, below. As
seen, the pre-treated mouse group was the highest in average
viability, the post-treated mouse group next and the untreated
mouse group is the lowest. The mice tested were not changed in
25 body weight.

TABLE 12

Immunotherapy of the Polysaccharide Substance

Test Groups (mg/kg)	Avg. Viability	Proportion (%)	Nos.of Viable Mice on the 30th day
Untreated	20.8	100	0/10
Pretreated (100)	29.0	139	8/10
Posttreated (100)	25.8	124	2/10

10 EXAMPLE XVII : Activity of the Polysaccharide Substance against
Cancer Cells according to Immunochemo Therapy

Experiment 1: Anticancer Activity according to Co-administration
of the Polysaccharide Substance and Adriamycin

15

A homogeneous plantation test system of B16F10 cutaneous
cancer cell was utilized for the measurement of the anticancer
activity upon co-administration of the polysaccharide substance
and adriamycin. The cutaneous cancer cells were implanted in the
20 peritoneal cavity of mice. Adriamycin was peritoneally
administered to the mice for 12 days after tumor implantation
while the polysaccharide substance was peritoneally administered
from a week before the implantation to 12 days after the
implantation. For 60 days, the viability of the mice was
25 recorded everyday. Adriamycin was used at such low amounts as
not to damage the mice.

The results are given as shown in Figs. 8a and 8b and summarized in Table 13, below. As apparent from the data, the mice which were not treated with the agents, were very poor in viability and the mice which were treated with the polysaccharide substance alone were far superior in viability to the untreated mice. In Figs. 8a and 8b, there are shown the results when adriamycin was administrated at a dose of 0.1 mg and 0.3 mg per kg of body weight of a mouse, respectively. The viability was much better at a dose of 0.3 mg/kg than at 0.1 mg/kg. In the case of co-administration with the polysaccharide substance, adriamycin also more raised the mouse viability at a dose of 0.3 mg/kg than at 0.1 mg/kg. Numerically, the mice which were implanted with B16F10 cutaneous cancer cells were alive for 18.1 days in average. When they were treated with the polysaccharide substance alone, their average viability was extended to 31 days. Adriamycin allowed the mice to be alive for 35.4 days in average when being singly administrated at a dose of 0.1 mg/kg while the mice could live 40.1 days in average by the sole administration of adriamycin at a dose of 0.3 mg/kg. Where adriamycin was administrated in combination with the polysaccharide substance, the mice's average viability was significantly extended from 46.9 days at a dose of 0.1 mg/kg to 57.8 days at a dose of 0.3 mg/kg. During the test, the mice did not lose body weight.

TABLE 13

Immunotherapy of Combinations of Adriamycin and the
Polysaccharide Substance

Test Group (mg/kg)	Avg. Viability	Proportion (%)	Nos. of viable mice on the 60th day
Untreated	18.1	100.0	0/10
P.S ¹ (100)	31.0	171.3	0/10
Am ² (0.1)	35.4	195.6	0/10
Am (0.1) + P.S. (100)	46.9	259.1	4/10
Am (0.3)	40.1	240.9	2/10
Am (0.3) + P.S (100)	57.8	319.3	9/10

¹Polysaccharide Substance

²Adriamycin

Experiment 2: Anticancer Activity according to Co-administration
of the Polysaccharide Substance and mitomycin C

A homogeneous plantation test system of B16F10 cutaneous
cancer cell was utilized for the measurement of the anticancer
activity upon co-administration of the polysaccharide substance
and mitomycin C, as in Experiment 1. The cutaneous cancer cells
were implanted in the peritoneal cavity of mice. Mitomycin C was
peritoneally administrated to the mice for 12 days after tumor
implantation while the polysaccharide substance was peritoneally
administrated from a week before the implantation to 12 days
after the implantation. For 60 days, the viability of the mice
was recorded everyday. Mitomycin C was used at such low amounts

as not to damage the mice.

Similar results to those of Experiment 1 were obtained and are summarized in Table 14, below. As apparent from the data, the mice which were not treated with the agents, were very poor in viability and the mice which were treated with the polysaccharide substance alone were far superior in viability to the untreated mice. Numerically, the mice which were implanted with B16F10 cutaneous cancer cells were alive for 18.1 days in average, but when they were treated with the polysaccharide substance alone, their average viability was extended to 31 days. mitomycin C allowed the mice to be alive for 32.5 days in average when being singly administrated at a dose of 0.2 mg/kg while the mice could live 38.0 days in average by the sole administration of mitomycin C at a dose of 0.6 mg/kg. Where mitomycin C was administrated in combination with the polysaccharide substance, the mice's average viability was significantly extended from 43.5 days at a dose of 0.2 mg/kg to 51.5 days at a dose of 0.6 mg/kg. During the test, the mice did not lose body weight.

TABLE 14

Immunotherapy of Combinations of Mitomycin C and the
Polysaccharide Substance

Test Group (mg/kg)	Avg. Viability	Proportion (%)	Nos. of viable mice on the 60th day
Untreated	18.1	100.0	0/10
P.S ¹ (100)	31.0	171.3	0/10
M.C ² (0.2)	32.5	179.6	0/10
M.C(0.2)+P.S(100)	43.5	240.3	3/10
M.C(0.6)	38.0	209.9	1/10
M.C(0.6)+P.S(100)	51.5	284.5	7/10

¹polysaccharide substance

²Mitomycin C

Experiment 3 : Anticancer Activity according to Co-administration
of the Polysaccharide Substance and Cisplatin

A homogeneous plantation test system of B16F10 cutaneous cancer cell was utilized for the measurement of the anticancer activity upon co-administration of the polysaccharide substance and cisplatin, as in the above experiments. The cutaneous cancer cells were implanted in the peritoneal cavity of mice. Cisplatin was peritoneally administrated to the mice for 12 days after tumor implantation while the polysaccharide substance was peritoneally administrated from a week before the implantation to 12 days after the implantation. For 60 days, the viability of the mice was recorded everyday. Cisplastin was used at such low

amounts as not to damage the mice.

Similar results to those of the above experiments were obtained and are summarized in Table 15, below. As apparent from the data, the mice which were not treated with the agents, were
5 very poor in viability and the mice which were treated with the polysaccharide substance alone were far superior in viability to the untreated mice. Numerically, the mice which were implanted with B16F10 cutaneous cancer cells were alive for 18.1 days in average, but when they were treated with the polysaccharide
10 substance alone, their average viability was extended to 31 days. Cisplatin allowed the mice to be alive for 33 days in average when being singly administrated at a dose of 0.5 mg/kg while the mice could live 40.5 days in average by the sole administration of cisplatin at a dose of 1.5 mg/kg. Where cisplatin was
15 administrated in combination with the polysaccharide substance, the mice's average viability was significantly extended from 44.0 days at a dose of 0.5 mg/kg to 52.6 days at a dose of 1.5 mg/kg. During the test, the mice did not lose body weight.

TABLE 15

Immunotherapy of Combinations of Cisplatin and the
Polysaccharide Substance

Test Group (mg/kg)	Avg. Viability	Proportion (%)	Nos. of viable mice on the 60th day
Untreated	18.1	100.0	0/10
P.S ¹ (100)	31.0	171.3	0/10
C.P ² (0.5)	33.0	182.3	0/10
C.P(0.5)+P.S(100)	44.0	243.0	3/10
C.P(1.5)	40.5	223.8	2/10
C.P(1.5)+P.S(100)	52.6	290.6	7/10

¹Polysaccharide Substance

²Cisplatin

Experiment 4 : Anticancer Activity according to Co-administration
of the Polysaccharide Substance and 5-fluoro uracil

A homogeneous plantation test system of B16F10 cutaneous cancer cell was utilized for the measurement of the anticancer activity upon co-administration of the polysaccharide substance and 5-fluoro uracil, as in the above experiments. The cutaneous cancer cells were implanted in the peritoneal cavity of mice. 5-fluoro uracil was peritoneally administrated to the mice for 12 days after tumor implantation while the polysaccharide substance was peritoneally administrated from a week before the implantation to 12 days after the implantation. For 60 days, the viability of the mice was recorded everyday. 5-fluoro uracil was

used at such low amounts as not to damage the mice.

Similar results to those of the above experiments were obtained and are summarized in Table 16, below. As apparent from the data, the mice which were not treated with the agents were very poor in viability and the mice which were treated with the polysaccharide substance alone were far superior in viability to the untreated mice. Numerically, 5-Fluoro uracil allowed the mice to be alive for 34.2 days in average when being singly administrated at a dose of 1 mg/kg and for 45.5 days in average when being administrated in combination with the polysaccharide substance. A dose of 3 mg/kg of 5-fluoro uracil enabled the mice to live 40.2 days in average when being administrated alone and to live 54.8 days in average when being administrated together with the polysaccharide substance. During the test, the mice did not lose body weight.

TABLE 16

Test Group (mg/kg)	Avg. Viability	Proportion (%)	Nos. of viable mice on the 60th day
Untreated	18.1	100.0	0/10
P.S ¹ (100)	31.0	171.3	0/10
5-FU ² (1)	33.0	187.8	0/10
5-FU(1)+P.S(100)	44.0	251.4	4/10
5-FU(3)	40.5	222	2/10
5-FU(3)+P.S(100)	52.6	302.0	8/10

¹ Polysaccharide Substance

² Fluorouracil

EXAMPLE XVIII : Activity of the Polysaccharide Substance against
Cancer Cells derived from the Human Body

The activity of the polysaccharide substance against human
5 cancer was assayed using NCl-H23, a lung cancer cell line. Nude
mice were subcutaneously implanted with NCl-H23 lung cancer
cells. The polysaccharide substance and adriamycin were
peritoneally administrated to the nude mice for 12 days after
cancer cell implantation. The size of a mass of the cancer cells
10 was measured and the cancer cells were separated and weighed on
the 19th day after cancer cell implantation. Because the nude
mice were devoid of T-cells, their immunological competency was
generally attributed to the activity of natural killer cells or
macrophage cells. Therefore, the immunological enhancement
15 obtained when administrating the polysaccharide substance was
mediated through these two type cells.

The results are given as shown in Figs. 9a and 9b and
numerically listed in Tables 17 and 18, below. As apparent from
these data, the cancer cells which were treated with neither
20 adriamycin nor the polysaccharide substance was the largest in
size, the polysaccharide substance-treated cancer cells next, and
the adriamycin-treated cancer cells the smallest. For weight,
the same order resulted in. The tested mice did not gain or lose
body weight.

TABLE 17

Immunotherapy for Human Cancer Cells

(Cell size mm³)

Test group	14th day	17th day	18th day	19th day
Untreated	43	105	131	158
P.S ¹ (100mg/kg)	20	39	40	43
Am ² (1mg/kg)	12	27	34	33

¹ Polysaccharide Substance² Adriamycin

TABLE 18

Immunotherapy for Human Cancer Cells

Test Group	Final Cancer Weight
untreated	259
P.S ¹ (100mg/kg)	113
Adriamycin (1mg/kg)	95

¹ Polysaccharide Substance

EXAMPLE XIX : Effect of the Polysaccharide Substance on

Metastasis of Mouse Cancer Cells

In order to test the polysaccharide substance for the prevention of cancer metastasis, B16F10 cutaneous cancer cells were implanted in the tail veins of C57BL/6 mice. On the 14th day after implantation, the lungs were isolated from the mice to investigate whether or to what extent cancer metastasis occurred.

Adriamycin was peritoneally administrated to the mice for 12 days after tumor implantation while the polysaccharide substance was peritoneally administrated from a week before the implantation to 12 days after the implantation. The B16F10 cutaneous cancer cells injected intravenously to the tails were translocated to form black spots on the lungs. The results are given as shown in Figs. 10a and 10b and numerically summarized in Table 19, below. As shown in Fig. 10a, as many as about 272 cancer spots were found in the mouse group which was not treated with any agents and they were reduced to 197 when administrating adriamycin at a dose of 1 mg/kg and further reduced to 172 when administrating adriamycin at a dose of 3 mg/kg. Adriamycin at a dose of 0.1-0.3 mg/kg showed a potent activity of preventing cancer cells from growing, but it showed only a very weak effect on the prevention of cancer metastasis event at a dose of 1-3 mg/kg. The number of the cancer spots was reduced to 83 by administrating the polysaccharide substance alone. 136 and 131 cancer spots were found in the mouse groups which were administrated with the polysaccharide substance in combination with adriamycin at a dose of 1 mg/kg and 3 mg/kg, respectively. Therefore, the polysaccharide substance is more effective for the prevention of cancer metastasis than adriamycin.

TABLE 19

Test Group	Nos. of Spots
Untreated	272
P.S ¹ (100mg/kg)	83
Adriamycin (1mg/kg)	197
Adriamycin (1mg/kg)+P.S (100mg/kg)	136
Adriamycin (3mg/kg)	172
Adriamycin (3mg/kg)+P.S (100mg/kg)	131

¹ Polysaccharide Substance

EXAMPLE XX : Test for Direct Cytotoxicity of the Polysaccharide Substance to Cancer Cells

The anticancer activity of the polysaccharide substance was tested for whether it is expressed through immunological enhancement or by direct cytotoxicity to cancer cells. B16F10 mouse cutaneous cancer cells and NCl-H23 human lung cancer cells were directly treated with the polysaccharide substance and adriamycin at an amount of 0.3-30 μ g/kg. The cancer cells still alive on the 2nd day after treatment were measured according to a sulforhodamin B method. The results are given as shown in Fig. 11 and numerically summarized in Table 20, below. The cell quantity of the test groups which were not treated with any chemicals was expressed as "100 %" and the cell quantities of treated test groups were compared with this standard. As apparent from the data, adriamycin exhibits potent cytotoxicity

to the two type cancer cells while the polysaccharide substance does not. Taken together, the data obtained in the above examples show that adriamycin exerts anticancer activity through cytotoxicity while the polysaccharide substance displays anticancer activity through the immunological enhancement.

TABLE 20

Direct Cytotoxicity of the Polysaccharide Substance

Test Group	Adriamycin		Polysaccharide Substance	
	B16F10	NC1-H23	B16F10	NC1-H23
Untreated	100	100	100	100
0.3 $\mu\text{g/mL}$	56	100	93	93
1 $\mu\text{g/mL}$	29	71	89	89
3 $\mu\text{g/mL}$	9	39	80	80
10 $\mu\text{g/mL}$	-10	7	78	78
30 $\mu\text{g/mL}$	-10	-1	77	77

EXAMPLE XXI : Test for Use as an Aid in Prophylaxis and Treatment of AIDS

The polysaccharide substance according to the present invention was tested for the availability of an aid for the prophylaxis and treatment of AIDS and found to be effective therefor.

Taken together, the data of the Examples show that the mushroom of the present invention is a novel strain with a

relatedness to *Phellinus* spp. as analyzed by RFLP and the polysaccharide substance obtained from the mycelia of *Phellinus* spp. has an anticancer activity by inciting the cytotoxicity of T-cells and the antibody forming potency of B-
5 cells. In addition, the polysaccharide substance of the present invention shows a potent activity against cancer metastasis. Consequently, the polysaccharide substance, which can stimulate the immunity of the body, may be effective in the prophylaxis and treatment of immune-related diseases, such as cancers and AIDS
10 and thus, is very useful in the biomedical industry.

The present invention has been described in an illustrative manner, and it is to be understood the terminology used is intended to be in the nature of description rather than of limitation.

15 Many modifications and variations of the present invention are possible in light of the above teachings. Therefore, it is to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described.